

91267



NEW ZEALAND QUALIFICATIONS AUTHORITY
MANA TOHU MĀTAURANGA O AOTEAROA

QUALIFY FOR THE FUTURE WORLD
KIA NOHO TAKATŪ KI TŌ ĀMUA AO!

2

SUPERVISOR'S USE ONLY

Level 2 Mathematics and Statistics, 2018

91267 Apply probability methods in solving problems

9.30 a.m. Wednesday 14 November 2018
Credits: Four

Achievement	Achievement with Merit	Achievement with Excellence
Apply probability methods in solving problems.	Apply probability methods, using relational thinking, in solving problems.	Apply probability methods, using extended abstract thinking, in solving problems.

Check that the National Student Number (NSN) on your admission slip is the same as the number at the top of this page.

You should attempt ALL the questions in this booklet.

Make sure that you have Formulae Sheet L2–MATHF.

Show ALL working.

If you need more space for any answer, use the page(s) provided at the back of this booklet and clearly number the question.

Check that this booklet has pages 2–12 in the correct order and that none of these pages is blank.

YOU MUST HAND THIS BOOKLET TO THE SUPERVISOR AT THE END OF THE EXAMINATION.

TOTAL

ASSESSOR'S USE ONLY

QUESTION ONE

Each day, temperatures are recorded every hour at various locations around New Zealand. The National Institute of Water and Atmospheric Research (NIWA) obtains the “daily temperature” by finding the mean of all the recorded temperatures in one 24-hour period at that location.

- (a) Over the last seven years, the daily temperatures in Reefton have been found to be approximately normally distributed with a mean of 11.6°C and standard deviation of 4.8°C .

Use this approximation to answer the questions in parts (i) to (iv).

Working and/or diagrams must be shown. Correct answer(s) alone will generally limit grades to Achievement.

- (i) Find the probability that a randomly chosen day in Reefton would have a daily temperature between 11.6°C and 18.8°C .

- (ii) Find the probability that a randomly chosen day in Reefton would have a daily temperature over 17°C .

- (iii) People in Reefton talk about the hottest 1% of days as being “scorchers”.

What is the lowest possible daily temperature of a “scorcher” day?

- (iv) Using this normal distribution approximation, the probability of a daily temperature lower than 9°C is 29.4% (3sf).

Suppose that temperatures in Reefton increased due to climate change, but the standard deviation remained the same.

What would the mean daily temperature for Reefton need to become to make this probability fall to 20.0% (3sf)?

QUESTION TWO

- (a) Matiu analysed Kaitaia's daily weather data from NIWA for the last seven years. He classified a day as "windy" if the mean wind speed was above 36 km/h, and "still" otherwise. He classified a day as "wet" if more than 2 mm of rain fell, and "dry" otherwise. He summarised his analysis in Table 1 below.

Table 1

Kaitaia	Wet	Dry	Total
Windy	553	1093	1646
Still	88	822	910
Total	641	1915	2556

Use Table 1 to answer the questions in parts (i) to (iii).

- (i) What proportion of days over this time were windy?

- (ii) What proportion of wet days over this time were still?

- (iii) Over these years, is it more likely to be wet on a windy day, or wet on a still day?
Support your answer with probability calculations.

QUESTION THREE

Nancy finds some data from NIWA on weather in Ashburton and in Timaru over the past seven years. She analyses the data and finds that:

- It was wet on 45% of days in Ashburton.
- If it was wet in Ashburton, the probability that it was wet on the same day in Timaru was 63%.
- If it was dry in Ashburton, the probability that it was dry on the same day in Timaru was 88%.

- (a) (i) Find the probability that it was dry in both Ashburton and Timaru on a randomly chosen day.

- (ii) Find the probability that, on a randomly chosen day, only one of the towns was wet.

- (iii) If it was a dry day in Timaru, what is the probability that it was also dry in Ashburton on the same day?

- (b) Nancy is planning a one-day bicycle ride from Ashburton, through Timaru and then to Waimate. Nancy's grandfather says that, in his experience, the probability of it being dry in all three towns on the same day is "somewhere between 30 and 35%".

On the day of Nancy's ride, it is dry in both Ashburton and Timaru.

If Nancy's grandfather is correct, what is the smallest probability that it is a wet day in Waimate?

(c) Nancy's friend Teri uses the NIWA data for the past seven years to find out that:

- 45% of days in Ashburton were wet
- 35% of days in Timaru were wet.

Teri constructs the tree diagram in Figure 1 below.

Figure 1




Figure 2



Adapted from: http://ortho.linz.govt.nz/nz_small_scale/si_1million.jpg

