

Assessment Schedule – 2015

Calculus: Apply the algebra of complex numbers in solving problems (91577)

Evidence

| Q1 | Expected Coverage | Achievement u | Merit r | Excellence t |
|-----|--|----------------------|-------------------|-----------------|
| (a) | $x = 4 \pm 2\sqrt{3}$ | Correct solution. | | |
| (b) | $u^3 = -8$ clearly marked on Argand diagram. | Correct solution. | | |
| (c) | $p(3 - 7i) + q(-4 + 6i) = 6.5 - 11i$ $3p - 7pi - 4q + 6iq = 6.5 - 11i$ $\Rightarrow 3p - 4q = 6.5$ and $-7p + 6q = -11$ $p = 0.5$ $q = -1.25$ | Correct second line. | Correct solution. | |
| (d) | $'b^2 - 4ac' = (2c + 1)^2 + 4 \times 3 \times (c + 3)$ $= 4c^2 + 16c + 37$ <i>OR</i> $= 4(c^2 + 4c) + 37$ $= (2c + 4)^2 + 21 \geq 0$ $= 4([c + 2]^2 - 4) + 37$ Since $(2c + 4)^2 \geq 0$ $= 4(c + 2)^2 + 21$ Which is always positive, as $(c + 2)^2$ is always positive. | Correct second line. | Correct solution. | |

| Q2 | Expected Coverage | Achievement u | Merit r | Excellence t |
|-----|--|-----------------------------------|---|-------------------|
| (a) | -23 | Correct solution. | | |
| (b) | $\frac{2+3i}{5+i} = \frac{(2+3i)(5-i)}{(5+i)(5-i)}$ $= \frac{13+13i}{26}$ $= \frac{13}{26}(1+i)$ $= \frac{1}{2}(1+i)$ $\therefore k = \frac{1}{2}$ | Correct solution. | | |
| (c) | $1 = A(x(x-1)) + B(x-1) + Cx^2$ $= Ax^2 - Ax + Bx - B + Cx^2$ $= (A+C)x^2 + (B-A)x - B$ $\therefore B = -1, A = -1, C = 1$ | Correct second line. | Correct solution. | |
| (d) | $\left(\frac{4i^7 - i}{1+2i}\right)^2 = \left(\frac{-4i - i}{1+2i}\right)^2$ $= \left(\frac{-5i}{1+2i}\right)^2$ $= \frac{-25}{-3+4i}$ $= \frac{-25(-3-4i)}{(-3+4i)(-3-4i)}$ $= \frac{75+100i}{25}$ $= 3+4i$ | Correct third line. Or CRO | Correct solution. | |
| (e) | $\frac{z-2}{z+5} = \frac{x+yi-2}{x+yi+5}$ $= \frac{(x-2)+yi}{(x+5)+yi}$ $= \frac{[(x-2)+yi][(x+5)-yi]}{[(x+5)+yi][(x+5)-yi]}$ $= \frac{x^2+y^2+3x-10+7yi}{x^2+10x+25+y^2}$ $\text{Arg} = \frac{\pi}{4} \Rightarrow x^2+y^2+3x-10=7y$ $\therefore x^2+y^2+3x-10-7y=0$ | Correct second line. | Correct 'simplification', 4th line. Accept correct factorised denominator. | Correct solution. |

| NØ | N1 | N2 | A3 | A4 | M5 | M6 | E7 | E8 |
|------------------------------------|---------------------|----|----|----|----|----|------------------------|-----|
| No response; no relevant evidence. | 1 partial solution. | 1u | 2u | 3u | 1r | 2r | 1t with 1 minor error. | 1 t |

| Q3 | Expected Coverage | Achievement u | Merit r | Excellence t |
|-----|--|--|---|-----------------|
| (a) | $zw = (4 + 2i)(-1 + 3i)$ $= -4 + 12i - 2i + 6i^2$ $= -10 + 10i$ $\text{Arg}(zw) = 135^\circ = \frac{3\pi}{4} = 2.36 \text{ rad}$ | Correct solution. | | |
| (b) | $b^2 - 4ac = 0$ $\frac{1}{k^2} - 4k \times 2 = 0$ $\frac{1}{k^2} = 8k$ $k^3 = \frac{1}{8}$ $k = \frac{1}{2}$ | Correct solution. | | |
| (c) | $f(-2) = 0$ $\Rightarrow -24 + 4A + 16 = 0$ $A = 2$ $3w^3 + Aw^2 - 3w + 10 = (w + 2)(3w^2 - 4w + 5)$ $\therefore (w + 2)(3w^2 - 4w + 5) = 0$ $w = \frac{4 \pm \sqrt{16 - 4 \times 3 \times 5}}{6}$ $= \frac{4 \pm \sqrt{-44}}{6}$ $= \frac{2 \pm \sqrt{11}i}{3}$ <p>Or $w = 0.667 \pm 1.1055i$</p> | Correct value of A found. | Correct solution. | |
| (d) | $z^3 = 2k \text{cis}\left(\frac{\pi}{3}\right) \quad \text{or Arg} = 1.047 \quad \text{or } 60^\circ$ <p>Therefore roots are:</p> $z_1 = \sqrt[3]{2k} \text{cis}\left(\frac{\pi}{9}\right) \quad \text{or Arg} = 0.349 \quad \text{or } 20^\circ$ $z_2 = \sqrt[3]{2k} \text{cis}\left(\frac{7\pi}{9}\right) \quad \text{or Arg} = 2.443 \quad \text{or } 140^\circ$ $z_3 = \sqrt[3]{2k} \text{cis}\left(\frac{13\pi}{9}\right) \quad \text{or Arg} = 4.538 \quad \text{or } 260^\circ$ <p>OR $\sqrt[3]{2k} \text{cis}\left(\frac{-5\pi}{9}\right) \quad \text{or Arg} = -1.745 \quad \text{or } -100^\circ$</p> | One correct solution or 3 correct arguments. | Three correct solutions – allow alternative correct values. | |

| | | |
|---|--|--------------------------|
| <p>(e)(i)</p> $z^5 = 1$ $z^5 = 1 \operatorname{cis} 0$ <p>Therefore roots are</p> $z_1 = 1 \operatorname{cis} 0$ <p>or 1</p> $z_2 = 1 \operatorname{cis} \left(\frac{2\pi}{5} \right) \quad \text{or Arg} = 1.257 \quad \text{or } 72^\circ$ <p>or $0.309 + 0.951i$</p> $z_3 = 1 \operatorname{cis} \left(\frac{4\pi}{5} \right) \quad \text{or Arg} = 2.513 \quad \text{or } 144^\circ$ <p>or $-0.809 + 0.588i$</p> $z_4 = 1 \operatorname{cis} \left(\frac{6\pi}{5} \right) \quad \text{or Arg} = 3.770 \quad \text{or } 216^\circ$ <p>or $1 \operatorname{cis} \left(\frac{-4\pi}{5} \right) \quad \text{or Arg} = -2.513 \quad \text{or } -144^\circ$</p> <p>or $-0.809 - 0.588i$</p> $z_5 = 1 \operatorname{cis} \left(\frac{8\pi}{5} \right) \quad \text{or Arg} = 5.027 \quad \text{or } 288^\circ$ <p>or $1 \operatorname{cis} \left(\frac{-2\pi}{5} \right) \quad \text{or Arg} = -1.257 \quad \text{or } -72^\circ$</p> <p>or $0.309 - 0.951i$</p> <p>(ii) Let $p = 1 \operatorname{cis} \left(\frac{2\pi}{5} \right)$ which is z_2 above or $p = 0.309 + 0.951i$</p> <p>Then</p> $p^2 = \left(1 \operatorname{cis} \left(\frac{2\pi}{5} \right) \right)^2 = 1 \operatorname{cis} \left(\frac{4\pi}{5} \right) = z_3$ $p^3 = \left(1 \operatorname{cis} \left(\frac{2\pi}{5} \right) \right)^3 = 1 \operatorname{cis} \left(\frac{6\pi}{5} \right) = z_4$ $p^4 = \left(1 \operatorname{cis} \left(\frac{2\pi}{5} \right) \right)^4 = 1 \operatorname{cis} \left(\frac{8\pi}{5} \right) = z_5$ <p>And $z_1 = 1 \operatorname{cis} 0 = 1$</p> <p>$\therefore$ roots are $1, p, p^2, p^3$ and p^4</p> | <p>Correct values of z_1 to z_5.</p> | <p>Correct solution.</p> |
|---|--|--------------------------|

| N0 | N1 | N2 | A3 | A4 | M5 | M6 | E7 | E8 |
|------------------------------------|---------------------|----|----|----|----|----|------------------------|-----|
| No response; no relevant evidence. | 1 partial solution. | 1u | 2u | 3u | 1r | 2r | 1t with 1 minor error. | 1 t |

Cut Scores

| Not Achieved | Achievement | Achievement with Merit | Achievement with Excellence |
|--------------|-------------|------------------------|-----------------------------|
| 0 – 6 | 7 – 12 | 13 – 18 | 19 – 24 |